

Preliminary estimating the contemporary sedimentation trend in dry valley bottoms of first-order catchments of different landscape zones of the Russian Plain using the ^{137}Cs as a chronomarker

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© Published under licence by IOP Publishing Ltd. A general trend of erosion processes over the last 50-60 years can be estimated by dating sediments washed off from arable lands and accumulated in the first-order dry valleys bottoms. Three small (first-order) catchments were chosen as objects of the study. They are located, respectively, in the southern part of the taiga zone, the zone of temperate broad-leaf forests and the forest-steppe zone of the Russian Plain. To date the sediments accumulated in the bottoms the radioactive caesium-137 (^{137}Cs) of global (since 1954) and Chernobyl origin (1986) had been used as a chronomarker. The average (for all the catchments) sedimentation rates during the global ^{137}Cs fallout period (1963(1954)-1986) are at least 0.88-2.71 cm per year. For the period that has passed since the Chernobyl accident (1986-2015(2016)) the average rates were 0.15-1.07 cm per year. The greatest reduction in the sedimentation rates is observed in the subzone of the southern taiga, the lowest one is in the forest-steppe zone of the Russian Plain. The main reason for such significant reduction in the rates of sedimentation of the soil erosion products in the dry valley bottoms was a reduction of surface runoff within the catchments during a snowmelt period, as well as crop-rotation changes there.

<http://dx.doi.org/10.1088/1755-1315/107/1/012022>

References

- [1] Porto P, Walling D E and Callegari G 2011 *Hydrological Processes* 25 900
- [2] Golosov V N, Panin A V and Markelov M V 1999 *Physics and Chemistry of the Earth, Part A: Solid Earth and Geodesy* 24 885
- [3] Parfitt R L, Baisden W T, Ross C W, Rosser B J, Schipper L A and Barry B 2013 *New Zealand Geoderma* 192 159
- [4] Zapata F 2002 *Handbook for the assessment of soil erosion and sedimentation using environmental radionuclides* 219 (Springer)
- [5] Golosov V N, Belyaev V R and Markelov M V 2013 *Hydrological Processes* 27 794
- [6] Golosov V N, Ivanova N N, Gusarov A V and Sharifullin A G 2017 *Eurasian Soil Science* 10 1208
- [7] Zhang X and Walling D E 2005 *Journal of Environmental Quality* 34 523
- [8] Silant'ev A N and Shkuratova I G 1983 *Detecting Industrial Soil Contaminants and Atmospheric Fallout against the Global Contamination Background* (Leningrad: Gidrometeoizdat)
- [9] Almgren S and Isaksson M 2006 *Journal of Environmental Radioactivity* 91 102
- [10] Karadeniz Ö, Çakir R and Karakurt H 2015 *Journal of Environmental Radioactivity* 146 34
- [11] Ciszewski D, Czajka A and Blazej S 2008 *Environmental Geology* 55 86

- [12] Kirchner G, Strebl F, Bossew P, Ehlken S and Gerzabek M H 2009 Journal of Environmental Radioactivity 100 720
- [13] Korobova E M, Dogadkin N N, Shiryaev A A, Kolotov V P, Kononkova N N and Turkov V A 2014 Journal of Geochemical Exploration 142 100
- [14] Golosov V, Gusarov A, Litvin L, Yermolaev O, Chizhikova N, Safina G and Kiryukhina Z 2017 Proceedings of the International Association of Hydrological Sciences 375 27